

Global Hydromorphone Consumption, 2009: (mg/capita)

Note: Countries not listed either did not report, or reported an amount of zero (0) for hydromorphone consumption, to the INCB for 2009

1	Canada	23.7483
2	Austria	7.7969
3	United States of America	7.0185
4	Germany	4.2818
5	Switzerland	2.8450
6	Slovenia	1.3099
7	Luxembourg	1.2263
8	Spain	1.0985
9	Czech Republic	0.9798
10	Sweden	0.8187
11	Ireland	0.7779
12	Belgium	0.6602
13	Hungary	0.6270
14	United Kingdom	0.5922
15	Denmark	0.5905
16	Republic of Korea	0.5367
17	Australia	0.5097
18	Slovakia	0.4562
19	Serbia	0.4089
20	Cayman Islands	0.3509
21	Italy	0.3337
22	France	0.2800
23	Iceland	0.2446
24	Netherlands	0.2405
25	Finland	0.1998
26	New Caledonia	0.1364
27	Portugal	0.1305
28	Croatia	0.0781
29	Norway	0.0763
30	Colombia	0.0483
31	Saudi Arabia	0.0475
32	Mexico	0.0289
33	United Arab Emirates	0.0280
34	Israel	0.0142
35	Malta	0.0098
36	Philippines	0.0086
37	Jordan	0.0040
38	Egypt	0.0033
39	Kuwait	0.0017
40	Venezuela	0.0013
41	Yemen	0.0012
42	China	0.0002

Pain & Policy Studies Group
University of Wisconsin
Carbone Cancer Center
WHO Collaborating Center